arXiv:1412.2671v1 [math.AT] 8 Dec 2014

On the (in)finiteness of the image of Reshetikhin-Turaev representations

Julien Korinman
Institut Fourier BP 74, UMR 5582
University of Grenoble I
38402 Saint-Martin-d'Hères cedex, France
email: julien.korinman@ujf-grenoble.fr

Abstract

We state a simple criterion to prove the infiniteness of the image of Reshetikhin-Turaev representations of the mapping class groups of surfaces at odd prime levels. We use it to study some of the Reshetikhin-Turaev representations associated to the one-holed torus and derive an alternative proof of the results of [6].

Keywords: Reshetikhin-Turaev representations, mapping class group, quantum representations, Topological Quantum Field Theory.

1 Introduction

At the end of the 80' Witten [15] constructed a $2+1$ dimensional TQFT using path integrals which gives a three-dimensional intepretation of the Jones polynomial and gives invariants for 3-dimensional closed oriented manifold equipped with some links and additional structure. Reshetikhin and Turaev ([12]) gave a rigorous construction using quantum groups and more recently Blanchet, Masbaum, Habegger and Vogel constructed these TQFTs by means of Kauffman bracket skein algebra ([1]) following the works of Lickorish ([9]). We will use the construction of the latter authors.

These TQFTs give rise to a tower of representations $\rho_p, p \geq 3$ of some central extensions of the mapping class group $\widetilde{\mathrm{Mod}(\Sigma_g)}$ of any closed oriented surface Σ, equipped with colored points, which act on finite rank modules $V_p(\Sigma)$. Here we will discuss whether these representations have finite image or not. Several studies have been made in that sense. The Reshetikhin-Turaev representations associated to a torus without marked points have finite image. This fact was known in the conformal field theory community (see [2] and references herein) and has been proved independently by Gilmer in [7]. In higher genus, the Reshetikhin-Turaev representations have finite image at level 3 and 6 (see [16]) and 4 (it is a character). In any other cases, they have infinite image. It was proved by Funar in [6] when $(g,p) \neq (2,10)$ (see also [10] for a proof that they contain an element of infinite order). The last remaining case $g=2$, $p=10$ was treated in [3]. Concerning the representation associated to a one holed torus, it was shown in [14] that they have infinite image for high enough level.

In this paper, we develop a simple criterion to check whether a given Reshetikhin-Turaev representation has infinite image or not. Unfortunately, this criterion only works when $p=r$ or $p=2r$ for r an odd prime. It permits us to recover the above results in these cases, and to study the finiteness of the representations associated to a one-holed torus, which could not be derived from previous papers.

Denote by ρ_p^c the level p Reshetikhin-Turaev representation associated to a torus $\mathcal{T}^c$ equipped with a band colored by $2c$. The main theorem is the following:

Theorem 1.1. *Let r be an odd prime and $p=r$ or $p=2r$. Then we have:*

1. *If* $2c = r - 3$*, then* ρ_{2r}^c *has finite image.*
2. *If* $c \equiv 1 \pmod 3$ *and* $r \neq 3, 5$*, then* ρ_{2r}^c *has infinite image.*
3. *If (*$c \equiv 3 \pmod 5$ *and* $r \equiv 2 \pmod 5$ *or* $r \equiv 3 \pmod 5$*) or (*$c \equiv 1 \pmod 5$ *and* $r \equiv 3 \pmod 5$*) or (*$c \equiv 2 \pmod 5$ *and* $r \equiv 2 \pmod 5$*), then* ρ_{2r}^c *has infinite image.*
4. *If* $c \leq \frac{r-7}{3}$*, then* ρ_{2r}^c *has infinite image.*

Acknowledgements: The author is thankful to his advisor Louis Funar for useful discussions. He acknowledge support from the grant ANR 2011 BS 0102001 ModGroup and the GEAR Network.

2 Complete positivity

Let Σ be a closed oriented surface with colored punctures and $p \geq 3$. The associated representations ρ_p act on a finite rank k_p-module $V_p(\Sigma)$ where $k_p = \mathbb{Z}[A, \frac{1}{p}, \kappa]$ quotiented by the relations $\phi_{2p}(A) = 0$, where ϕ_{2p} represents the $2p$-th cyclotomic polynomial, and $\kappa^6 = A^{-6-\frac{p(p+1)}{2}}$. A choice of a particular $2p$-th primitive root of unity A and a compatible complex $\kappa \in \mathbb{C}$ give a complex vector space $V_p^{A,\kappa}(\Sigma) = V_p(\Sigma) \otimes \mathbb{C}$ and a complex representation $\rho_p^{A,\kappa}$ whose action is preserved by a non-degenerate Hermitian form $\langle \cdot, \cdot \rangle_p^{A,\kappa}$. Note that in [1], the invariant pairing is bilinear whereas in this paper we choose it Hermitian using the involution of k_p sending A to A^{-1}.

Definition 2.1. We say that $V_p(\Sigma)$ is *completely positive* if for any choice of $A \in \mathbb{C}$ as a primitive $2p$-th root of unity and compatible $\kappa \in \mathbb{C}$, the Hermitian form $(V_p^{A,\kappa}(\Sigma), \langle \cdot, \cdot \rangle_p^{A,\kappa})$ is positive definite or negative definite.

The only effect of changing κ is possibly to change the eigenvalues of the Hermitian form to its opposite so complete positivity only means that all eigenvalues have the same sign for a given κ and all A.

Proposition 2.2. *Let* r *be an odd prime number and* $p = r$ *or* $p = 2r$*. Then* $(V_p(\Sigma), \langle \cdot, \cdot \rangle_p)$ *is completely positive if and only if* ρ_p *has finite image.*

If $\rho_p(\widetilde{\mathrm{Mod}(\Sigma_g)})$ is finite, the following lemma, together with Roberts' results ([13]), which states that ρ_p is irreducible at prime levels, imply that $V_p(\Sigma)$ must be completely positive.

Lemma 2.3. *Let* $\rho : G \to \mathrm{GL}(V)$ *be an irreducible group representation on a finite dimensional complex vector space* V *equipped with a non degenerate invariant Hermitian form* $\langle \cdot, \cdot \rangle_0$*. If* $\rho(G)$ *is finite, then* $\langle \cdot, \cdot \rangle_0$ *is definite either positive or negative.*

Proof. Let $\langle , \rangle$ denotes an arbitrary scalar product on V. If $\rho(G)$ is finite, the Hermitian form on V defined by

$$\langle u, v \rangle_1 := \frac{1}{\#\rho(G)} \sum_{\rho(g) \in \rho(G)} \langle \rho(g)u, \rho(g)v \rangle$$

is definite positive and invariant under the action of G.

If $t \in [0, 1]$, we define the invariant Hermitian form

$$\langle u, v \rangle_t := t \langle u, v \rangle_0 + (1 - t) \langle u, v \rangle_1$$

Suppose there exists $v_0 \in V$ so that $\langle v_0, v_0 \rangle_0 < 0$. Since $\langle v_0, v_0 \rangle_1 > 0$, there exists $t_0 \in (0, 1)$ so that $\langle v_0, v_0 \rangle_{t_0} = 0$. The invariance of $\langle , \rangle_{t_0}$ under the action of G, implies that $\langle \rho(g)v_0, \rho(g)v_0 \rangle_{t_0} = 0$ for all $g \in G$. Since ρ is irreducible, the vector v_0 is cyclic so $\langle \cdot, \cdot \rangle_{t_0} = 0$ and $\langle \cdot, \cdot \rangle_0 = \frac{t_0 - 1}{t_0} \langle \cdot, \cdot \rangle_1$ is definite negative. □

When $p = r$ or $p = 2r$ with r an odd prime,
we write $\alpha_p := \begin{cases} p & \text{, if } p \equiv 3 \pmod 4 \\ 4r & \text{, if } p \equiv 1, 2 \pmod 4 \end{cases}$ and $\mathcal{O}_p := \mathbb{Z}[A] \Big/ \phi_{\alpha_p}(A)$.

It was showed in [8, 11] that $V_p(\Sigma)$ contains a free $\mathcal{O}_p$ lattice of maximal rank preserved by $\widetilde{\mathrm{Mod}(\Sigma_g)}$.

We denote by $\mu(\alpha_p) = \{q_1, \ldots, q_{\phi(\alpha_p)}\}$ the set of primitive α_p-th roots of unity and by $\phi(\alpha_p) = \#\mu(\alpha_p)$ the Euler totient function.

Lemma 2.4. *Let $p = r$ or $p = 2r$ with r an odd prime. Then the injective linear map*

$$\Psi : \mathcal{O}_p \to \mathbb{C}^{\phi(\alpha_p)}$$

sending A^n to $(q_1^n, \ldots, q_{\alpha_p}^n)$, has a discrete image.

Proof. We endow $\mathbb{C}^{\phi(\alpha_p)}$ with the norm:

$$\left\|(z_1, \ldots, z_{\phi(\alpha_p)})\right\|^2 := \frac{1}{\phi(\alpha_p)} \sum_i |z_i|^2$$

where $|z|^2$ represents the Euclidean norm in $\mathbb{C}$.

If $P(X) = \sum_i n_i X^i \in \mathcal{O}_p$, then:

$$\begin{aligned} \|\Psi(P)\|^2 &= \frac{1}{\phi(\alpha_p)} \sum_{q \in \mu(\alpha_p)} \sum_{i,j} n_i n_j q^{i-j} \\ &= \sum_i n_i^2 + \sum_{i \neq j} n_i n_j \left(\frac{1}{\phi(\alpha_p)} \sum_{q \in \mu(\alpha_p)} q^{i-j} \right) \\ &= \begin{cases} \sum_i n_i^2 & \text{, if } p \text{ is prime.} \\ \sum_i n_i^2 - 2 \sum_{|i-j|=2r} n_i n_j & \text{, if } p = 2r. \end{cases} \end{aligned}$$

Thus we have $\|\Psi(\mathcal{O}_p)\|^2 \in \mathbb{N}$, so $\Psi(\mathcal{O}_p)$ is discrete in $\mathbb{C}^{\phi(\alpha_p)}$. □

Proof of Proposition 2.2. If $\rho_p(\widetilde{\mathrm{Mod}(\Sigma_g)})$ is finite, then the irreducibility of ρ_p ([13]) together with Lemma 2.3, imply that every invariant forms $\langle \cdot, \cdot \rangle_p^{A,\kappa}$ must be either positive or negative, so $V_p(\Sigma)$ is completely positive.

Conversely, if $V_p(\Sigma)$ is completely positive, using the $\mathcal{O}_p$ lattice of [8, 11], we have an injective homomorphism from $\rho_p(G)$ to the group of matrices with coefficients in $\mathcal{O}_p$. Once composed with the map Ψ of Lemma 2.4, we get an injective group morphism $\widetilde{\Psi} : \rho(G) \to \mathrm{GL}_d(\mathbb{C}) \times \ldots \times \mathrm{GL}_d(\mathbb{C})$, where $d := dim(V_p(\Sigma))$. Lemma 2.4 implies that $\widetilde{\Psi}$ has a discrete image, and the complete positivity of $V_p(\Sigma)$ implies that the image lies in the compact product of $r - 1$ unitary groups. This implies the finiteness of $\rho_p(\widetilde{\mathrm{Mod}(\Sigma_g)})$. □

3 (In)finiteness of the one holed torus representations

Using Proposition 2.2, we can prove the main theorem of this paper:

(Proof of Theorem 1.1). For $0 \leq i \leq r-2-2c$, denote by $u_i^c \in V_p(\mathcal{T}^c)$ the vector associated to a lollipop graph whose stick is colored by $2c$ and loop colored by $i+c$, that is:

$u_i^c :=$ 

Using Theorem 4.11 of [1], and using the usual quantum numbers defined by $[n] := \frac{A^{2n}-A^{-2n}}{A^2-A^{-2}} \in k_p$, we have:

$$\begin{aligned} \frac{\langle u_{i+1}^c, u_{i+1}^c \rangle_p}{\langle u_i^c, u_i^c \rangle_p} &= \frac{[2c+i+2][i+1]}{[c+i+2][c+i+1]} \\ \frac{\langle u_{i+2}^c, u_{i+2}^c \rangle_p}{\langle u_i^c, u_i^c \rangle_p} &= \frac{[2c+i+3][2c+i+2][i+2][i+1]}{[c+i+1][c+i+3][c+i+2]^2} \end{aligned}$$

Proposition 2.2 states that ρ_p^c has infinite image if and only if there exists a primitive $2p$-th root of unity A such that one of the above numbers is negative.

When $2c = r-3$, choose $A = \exp\left(\frac{i\pi k}{r}\right)$ with $g.c.d.(k, 2r) = 1$. The vectors $\{u_0^c, u_1^c\}$ form a basis of $V_p(\mathcal{T}^c)$ and we have:

$$\frac{\langle u_1^c, u_1^c \rangle_p}{\langle u_0^c, u_0^c \rangle_p} = \frac{\sin(\frac{r-1}{r}\pi k)\sin(\frac{\pi k}{r})}{\sin(\frac{r+1}{2r}\pi k)\sin(\frac{r-1}{2r}\pi k)}$$

This product appears to be always positive for every k: indeed $\sin(\frac{r-1}{r}\pi k)$ and $\sin(\frac{\pi k}{r})$ are both positive if $k < r$ and both negative elsewhere whereas $\sin(\frac{r+1}{2r}\pi k)$ and $\sin(\frac{r-1}{2r}\pi k)$ are both positive if $\frac{k-1}{2}$ is even and both negative elsewhere. Thus ρ_p^c has finite image.

When $c \equiv 1 \pmod 3$, we take $k = \begin{cases} \frac{2r+1}{3} & \text{, if } r \equiv 1 \pmod 3 \\ \frac{2r-1}{3} & \text{, if } r \equiv 2 \pmod 3 \end{cases}$ and we have $\frac{\langle u_2^c, u_2^c \rangle_p}{\langle u_0^c, u_0^c \rangle_p} < 0$.

If $r \equiv 2 \pmod 5$, we put $k = \frac{2r+1}{5}$ and we have $\frac{\langle u_2^c, u_2^c \rangle_p}{\langle u_0^c, u_0^c \rangle_p} < 0$.

If $r \equiv 3 \pmod 5$, we put $k = \frac{2r-1}{5}$ and we have $\frac{\langle u_2^c, u_2^c \rangle_p}{\langle u_0^c, u_0^c \rangle_p} < 0$ when $c \equiv 1, 3 \pmod 5$.

When $c = 1$, then $\frac{\langle u_3^c, u_3^c \rangle_p}{\langle u_0^c, u_0^c \rangle_p} = \frac{[6]}{[2]} = \frac{\sin\left(\frac{2\pi k}{r}\right)}{\sin\left(\frac{4\pi k}{r}\right)}$ is negative if $\frac{r}{12} < k < \frac{r}{6}$. Such a k exists when $r \geq 12$.

For $r = 1$ or 7, we take $k = 1$ and for $r = 13, 17$, we take $k = 2$, and find again $\frac{\langle u_3^c, u_3^c \rangle_p}{\langle u_0^c, u_0^c \rangle_p} < 0$.

Finally, note that if there exist $0 \leq i \leq r-3$ and an odd $k < r$ so that $\frac{r}{2c+i+2} < k < \frac{r}{c+i+2}$ then $\frac{\langle u_{i+1}^c, u_{i+1}^c \rangle_p}{\langle u_i^c, u_i^c \rangle_p} < 0$ for $A = \exp\left(\frac{i\pi k}{r}\right)$. Then we can find such a i for $k = 3$ whenever $3 \in \bigcup_{i=0}^{r-3-2c} [\frac{r}{2c+i+2}, \frac{r}{c+i+2}] = [\frac{r}{r-1}, \frac{r}{c+2}]$, i.e. when $c \leq \frac{r-7}{3}$. □

Remark. We could have recovered the finiteness in the case where $2c = r - 3$ by using a theorem of Formanek (see [4]) which states that the only 2-dimensional irreducible representations of $B_3 = \langle t, t' | tt't = t'tt' \rangle$ are conjugate to one of the $\chi(y) \otimes \beta_3(q)$, where $\chi(y)$ is the character which sends both generators t and t' to $y \in \mathbb{C}^*$ and $\beta_3(q)$ is the reduced Burau representation in q. Since the eigenvalues of $\rho_p(t)$ are $\mu_c = (-1)^c A^{c(c+2)}$ and $\mu_{c+1} = (-1)^{c+1} A^{(c+1)(c+3)}$ we must have $y = -\mu_c$ and $q = A^r$ or $y = -\mu_{c+1}$ and $q = A^{-r}$ so q is a $4 - th$ primitive root of unity and $\beta_3(q)$ appears to have finite image (see [5] for a detailed discussion of finiteness of reduced Burau representations at roots of unity).

3.1 (In)finiteness of Reshetikhin-Turaev representations associated to closed surfaces

The following theorem results from [7, 6, 16, 3]. We derive another proof from Proposition 2.2.

Theorem 3.1 (*[7, 6, 16, 3]*)**.** *Let $g \geq 1$ and $p = r$ or $p = 2r$ with r an odd prime. Then $\rho_{p,g}$ has finite image if and only if $g = 1$ or $r = 3$.*

Proof. If $g = 1$, then $\frac{\langle u_{i+1}, u_{i+1} \rangle_p}{\langle u_i, u_i \rangle_p} = 1$ for all i, so $\rho_{p,1}$ is completely positive.

If $p = 3$ then $V_3(\Sigma_g)$ is one dimensional so is completely positive. If $p = 6$, we have $I_6 = \{0, 1\}$ and $(1, 1, 1)$ is not 6-admissible so, if Γ is a trivalent graph and σ a coloration of Γ, then u_σ is a union of disjoint circles, say n ones, colored by 1. Thus $\langle u_\sigma, u_\sigma \rangle_6 = 1$ and $V_6(\Sigma_g)$ is completely positive.

Now consider the case when $r = 5$ and $g = 2$. Take $\Gamma =$  and consider the associated vectors $u_{a,b,c}$ for (a, b, c) a p-admissible triple.

If $p = 5$, then $\frac{\langle u_{2,2,2}, u_{2,2,2} \rangle_5}{\langle u_{0,0,0}, u_{0,0,0} \rangle_5} = \frac{[4]}{[2]^2 [3]^2}$ has the sign of $[4]$ which is negative for $A = \exp\left(\frac{3i\pi}{5}\right)$.

If $p = 10$, then $\frac{\langle u_{2,1,1}, u_{2,1,1} \rangle_{10}}{\langle u_{0,0,0}, u_{0,0,0} \rangle_{10}} = \frac{\langle 2,1,1 \rangle}{\langle 1 \rangle^2 \langle 2 \rangle}$ has the sign of $\langle 2 \rangle = [3]$ which is negative for $A = \exp\left(\frac{3i\pi}{10}\right)$.

Now the fact that $V_p(\Sigma_2)$ is not completely positive for $r = 5$ implies that $V_p(\Sigma_g)$ is not completely positive when $g \geq 3$ as well, for we can embed the Theta graph into a genus $g \geq 3$ graph and define vectors $u_{a,b,c} \in V_p(\Sigma_g)$, with same norm as in genus 2, by coloring by 0 the complementary of the embedding.

Finally, if $r \neq 3, 5$ and $g \geq 2$, we isolate one handle of Σ_g and write $\Sigma_g \cong \Sigma_1 \bigcup_{S^1} \Sigma_{g-1}$. Then using Theorem 1.14 of [1], we have:

$$V_p(\Sigma_g) \cong \oplus_{c \in I_p} V_p(\mathcal{T}^c) \otimes V_p(\Sigma_{g-1}^c)$$

So the fact that $V_p(\mathcal{T}^1)$ is not completely positive (Theorem 1.1) implies that $V_p(\Sigma_g)$ is not completely positive either for $g \geq 2$. We conclude using Proposition 2.2. □

References

[1] C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel. Topological quantum field theories derived from the Kauffman bracket. *Topology*, 34(4):883–927, 1995.

[2] A. Coste and T. Gannon. Congruence subgroups and rational conformal field theory. *ArXiv Mathematics e-prints*, September 1999.

[3] J. K. Egsgaard and S. Fuglede Jørgensen. The homological content of the Jones representations at $q = -1$. *ArXiv e-prints*, February 2014.

[4] E. Formanek. Braid group representations of low degree. *Proc. London Math. Soc. (3)*, 73(2):279–322, 1996.

[5] M. Freedman and V. Krushkal. On the asymptotics of quantum SU(2) representations of mapping class groups. *Forum Math.*, 18(2):293–304, 2006.

[6] L. Funar. On the TQFT representations of the mapping class groups. *Pacific J. Math.*, 188(2):251–274, 1999.

[7] P.M. Gilmer. On the Witten-Reshetikhin-Turaev representations of mapping class groups. *Proc. Amer. Math. Soc.*, 127(8):2483–2488, 1999.

[8] P.M. Gilmer and G. Masbaum. Integral lattices in TQFT. *Ann. Sci. École Norm. Sup. (4)*, 40(5):815–844, 2007.

[9] W. B. R. Lickorish. Invariants for 3-manifolds from the combinatorics of the Jones polynomial. *Pacific J. Math.*, 149(2):337–347, 1991.

[10] G. Masbaum. An element of infinite order in TQFT-representations of mapping class groups. In *Low-dimensional topology (Funchal, 1998)*, volume 233 of *Contemp. Math.*, pages 137–139. Amer. Math. Soc., Providence, RI, 1999.

[11] K. Qazaqzeh. Integral lattices of the $SU(2)$-TQFT-modules. *Kobe J. Math.*, 30(1-2):19–32, 2013.

[12] N. Reshetikhin and V. G. Turaev. Invariants of 3-manifolds via link polynomials and quantum groups. *Invent. Math.*, 103(3):547–597, 1991.

[13] J. Roberts. Irreducibility of some quantum representations of mapping class groups. *J. Knot Theory Ramifications*, 10(5):763–767, 2001. Knots in Hellas '98, Vol. 3 (Delphi).

[14] R. Santharoubane. Limits of the quantum SO(3) representations for the one-holed torus. *ArXiv e-prints*, February 2012.

[15] E. Witten. Quantum field theory and the Jones polynomial. *Comm. Math. Phys.*, 121(3):351–399, 1989.

[16] G. Wright. The Reshetikhin-Turaev representation of the mapping class group at the sixth root of unity. *J. Knot Theory Ramifications*, 5(5):721–739, 1996.